

A large, semi-transparent blue graphic with rounded corners covers the left side of the page. Inside this graphic is a faded image of a laboratory setting, showing a person wearing gloves using a pipette to transfer liquid into a multi-well plate.

PCR CLIENT SERVICES MANUAL

WELCOME LETTER

Dear Customer,

Thank you for trusting Streamline Scientific for your molecular, microbiology, and toxicology testing. As a physician-founded company, we are passionate about the services we provide and know that the appropriate diagnostic information informs your decision making, and improves patient care.

Since our start in 2011, we have focused on simplifying the delivery of molecular technologies for both medical practices and patients. We do this by arming clinicians with the insights needed to prescribe the most appropriate treatment - within 24 hours.

Our sister organization, Molecular Designs, is a leading developer and manufacturer of PCR assays. In March of 2020, upon the news of the novel coronavirus, Molecular Designs was one of the first of 5 labs in the nation to submit an EUA and be granted FDA authorization.

This spirit of innovation continues today, developing novel tests to meet market needs and exploring new ways to better serve our many Alabama-based customers. Several of the elements that define the "Streamline Difference" include:

- Industry-leading technology and customer portal
- EMR integrations
- Unmatched customer support and dedicated team
- Door-to-door courier service
- An improved patient experience with multiple tests from a single swab

You are in exceptional hands with our Business Development and Client Relations teams, and the enclosed documentation will serve as a useful resource.

Should you have any questions or input on how we can better serve you, please don't hesitate to contact me directly.

Sincerely yours,



SHAWN HOOD

President

shood@StreamlineSci.com

DIRECTORY



BIRMINGHAM

2868 Acton Road
Suite 207
Birmingham, AL 35243

CLIA – 01D2074949
COLA – 25348



AUSTIN

13413 Galleria Circle
Suite Q-140
Bee Cave, TX 78738

CLIA – 45D2268534
COLA – 32244

CLIENT SERVICES CONTACT INFORMATION

Monday – Sunday: 8AM – 5PM

Email: clientservices@streamlinesci.com

Toll Free: 855.319.4459

Fax: 877.796.6185

BILLING SERVICES CONTACT INFORMATION

Email: patientbilling@streamlinesci.com

Local: 855.319.4459

LAB OPERATIONS

7 days a week

(Any holiday closures will be announced in advance and posted on streamlinesci.com)

NEW CLIENT SERVICES PORTAL

Easily order supplies, schedule pickups, or get help with any special requests from our dedicated account managers.

USERNAME: _____

PASSWORD: _____

COURIER BOX CODE: _____

BEGIN AT: →

[StreamlineSci.com](https://www.streamlinesci.com)



CREATE A TICKET

Customers may create a client services ticket from the home portal screen under the main menu option.

The screenshot shows a green header bar with the text "Client Services Request". Below the header, there are five buttons: "Request Supplies", "Request Pickup", "Create Ticket", "My Past Requests", and "Cancel". The "Create Ticket" button is highlighted with a yellow border.

Under Client Services, select "Create Ticket"

[Create Ticket](#)



SPECIAL REQUESTS, RESOLVED

The screenshot shows a "Request Pickup" form. The "Client" field is set to "Assurance Pathology Laboratory". The "Route" field is set to "Unassigned". The "Scheduled For (CST)" field is set to "04 / 25 / 2022" at "2:01 PM". There are checkboxes for "STAT" and "Complete". The "Notes" field contains the text "specimens. STAT pickup needed.". At the bottom, there are "Save" and "Cancel" buttons.

Our team is here to help with anything you need.

Make a special request

Use the "Link" button to **link your ticket to an item** within our lab information system—such as an existing lab order, a clinic location, provider, or patient profile—or to an uploaded document.

[Link](#)



SUPPLIES SIMPLIFIED

From collection kits and forms to transport labels, we have what you need to safely and efficiently manage your testing program.

Order supplies

- Select a **clinic location** and **contact**
- **Enter a description** for the request
- Select **needed supplies** from the drop-down menu of options
- **Enter a quantity** for each
- Click **“Save”** to submit your request

Save



SCHEDULE YOUR PICKUP TIME

It's never been easier to schedule a date and time for pickup. Enter your information along with any special requests and we'll take care of the rest.

Schedule a pickup

- Click the **“Now”** button to **autofill with the current date and time**

Now

- Use the Notes field to **enter any applicable details**, such as confirming the number of specimens to be picked up, or alerting the lab that this is a STAT request.
- Click **“Save”** to complete the pickup request.

Save

Visit the New Client Services Portal today to get started

StreamlineSci.com

SAMPLE COLLECTION GUIDELINES

COLLECTION CONTAINERS

- Synthetic Flocked Swab with Liquid Amies Transport buffer.
- Urine Vacutainers - Urine culture specimens only.
- Sterile Toxicology Cup with temperature strip (i.e., Clicktainer Vial, Temperature Strip Label) – Toxicology Samples Only.
- Sterile Dry Tubes (i.e., Transport Tube, 5ml, Capped, Self-Standing, Sterile) – Fungal Infection Panel ONLY.

SPECIMEN TYPE CONSIDERATIONS

PCR Testing:

- Swabs – Ideal samples are taken from the source of the potential infection. Ex. Nasopharyngeal, oropharyngeal, throat, superficial/subdermal wounds, vaginal tract, etc.
- Urine – If sending in urine specimens, **clean-catch urine** is required for UTI. Clean-catch urine is **not recommended** for STI or Candida testing.

Antimicrobial Susceptibility Testing:

- Swabs - Antimicrobial susceptibility testing is not recommended for nasal swabs due to the lack of clinical significance.
- Urine (bacteriostatic preservative) – No special requirements.

REJECTION CRITERIA

General criteria (applicable to all tests and sample types):

- Specimen is unlabeled or improperly labeled. Properly labeled includes two patient identifiers matching submitted requisition form.
- Requisition form is missing or incomplete.
- Specimen is improperly sealed, resulting in leakage and possible contamination during transport.
- Incorrect specimen type for selected panel.
- Insufficient sample volume for testing.

Swab Specimens:

- Swab specimens received with no original swab present in tube.
- Dry swab with no buffer present in tube.

Urine Specimens for UTI Testing:

- Urine specimens submitted in Aptima buffer or other container with no bacteriostatic preservative.

PCR Testing (any panel):

- Specimens received 6 days post-collection.

Culturing and Antimicrobial Susceptibility Testing:

- Swab specimens received 3 days post-collection.
- Swab specimens received in a non-liquid Amies buffer (such as viral transport medium).
- Nasal swab specimens (exceptions – see specimen type considerations).
- Urine specimens received 3 days post-collection.
- STI or Vaginitis specimens – Antimicrobial Susceptibility Testing is not available.

GENERAL DISCLAIMERS

Applies to all patient specimens submitted to Streamline Scientific: Proper patient identification and accurate specimen labeling is required per CLIA (Clinical Laboratory Improvement Amendment). Streamline Scientific has determined the analytical performance characteristics of the molecular tests. They have not been cleared or approved by the U.S. Food and Drug Administration. It is preferable that all specimen types are received within 2 days of collection. They will need to be received within the specimen stability range stated by the manufacturer of the collection device measured from the time of collection to time of receipt for laboratory testing, to prevent a disclaimer on the test report.

1 Patient Information

Last Name / First Name /

Address /

City / State / Zip

Phone

DOB

SSN

Insurance

Subscriber ID

Group #

Bill to: ☐ Insurance ☐ Facility
☐ Uninsured☐ Male
☐ Female

Race:

☐ Asian
☐ Black
☐ Caucasian
☐ Hispanic
☐ Native American
☐ Other
☐ N/A

Ethnicity:

☐ Hispanic
☐ Non-Hispanic
☐ N/A

2 Provider Information

Client Name / Account

Address / APT#

City / State / Zip

Phone

Ordering Provider

Collection
Date

Specimen Collected

State Collected

Collection
Time

3 Medical Necessity

As part of my antibiotic stewardship policy, I find it medically necessary to rapidly determine a patient's antibiotic resistance in order to treat with the most effective and timely data available to me direct to the patient. I hereby authorize Streamline Scientific to use (50% according to the CDC) for all co-pays and out of network charges.

Provider Signature

☐ Verbal Order
☐ Standing Order

4 Provider Information

The information I have provided on this form is accurate. I authorize Streamline Scientific to release this information to the patient or their family. I hereby authorize Streamline Scientific to use (50% according to the CDC) for all co-pays and out of network charges.

Patient Signature

Date

5 Panel List

☐ COVID-19 Only
☐ COVID/Flu/RSV☐ COVID-19 (Coronavirus)
☐ Influenza A & B
☐ RSV (types A & B)☐ COVID Respiratory Lite
(includes all the pathogens in the panel)☐ Haemophilus influenzae
☐ Moraxella catarrhalis
☐ Mycoplasma pneumoniae
☐ Strep. pyogenes (Group A)☐ COVID Respiratory
(includes all the pathogens in the panel)☐ Adenovirus
☐ Bocavirus
☐ Bordetella pertussis
☐ Chlamydia pneumoniae
☐ Coronavirus (229E, HKU1, NL63, OC43)
☐ EBV (mononucleosis)
☐ Enterovirus
☐ HMPV A & B
☐ Parainfluenza virus (type 1-4)
☐ Rhinovirus (types A & B)
☐ Staphylococcus aureus
☐ Streptococcus pneumoniae☐ COVID Respiratory Plus
(includes all the pathogens in the panel)☐ Acinetobacter baumannii
☐ Enterobacter cloacae
☐ Klebsiella aerogenes
☐ Klebsiella pneumoniae
☐ Legionella pneumophila
☐ Proteus mirabilis
☐ Pseudomonas aeruginosa
☐ Staphylococcus epidermidis
☐ ABX Resistance Marker
Methicillin/Oxacillin (mecA)

ICD 10 CODES

☐ R09.81 Congestion
☐ R50.9 Fever
☐ J02.9 Pharyngitis
☐ Z20.89 Exposure
☐ R05.9 Cough, unspecified

COVID VACCINATION STATUS:

☐ Z28.311 Partially vaccinated for COVID-19
☐ Z28.310 Unvaccinated for COVID-19
☐ Z28.39 Other immunization status

SPECIMEN SOURCE

☐ Nasal Swab
☐ Ear Swab
☐ Nasopharyngeal Swab
☐ Oropharyngeal Swab
☐ Sputum
☐ Other: _____☐ UTI w/ ABX Resistance☐ Acinetobacter baumannii
☐ Bacteroides fragilis
☐ Citrobacter braakii/freundii
☐ Citrobacter koseri
☐ Enterobacter cloacae
☐ Enterococcus spp.
☐ Escherichia coli
☐ Klebsiella aerogenes
☐ K. oxytoca/michiganensis
☐ Klebsiella pneumoniae
☐ Morganella morganii
☐ Proteus mirabilis
☐ Pseudomonas aeruginosa
☐ Serratia marcescens
☐ Staphylococcus aureus
☐ Staphylococcus epidermidis
☐ Staphylococcus saprophyticus
☐ Strep. pyogenes (Group A)

ABX Resistance Markers

☐ β -lactamase (blaKPC)
☐ β -lactamase (CTX-M-Group 1)
☐ metallo- β -lactamase (blaNDM)
☐ Fluoroquinolones
☐ Methicillin/Oxacillin (mecA)
☐ Sulfonamides
☐ Trimethoprim☐ UTI Plus
(includes all the pathogens in the panel)☐ Candida albicans
☐ Candida dubliniensis
☐ Candida glabrata
☐ Candida krusei
☐ Candida parapsilosis
☐ Candida tropicalis
☐ Mycoplasma genitalium
☐ Mycoplasma hominis
☐ Prevotella bivia
☐ Strep. agalactiae (Group B)
☐ Ureaplasma urealyticum

ICD 10 CODES

☐ R35.0 Frequency of Micturition
☐ Z22.39 Carrier of other specified bacterial disease
☐ R30.0 Dysuria
☐ N30.00 Acute cystitis without hematuria
☐ N30.20 Other chronic cystitis w/o hematuria
☐ N41.0 Acute prostatitis

SPECIMEN SOURCE

☐ Clean catch urine
☐ Urethral swab☐ Wound/Derm w/ ABX Resistance☐ Acinetobacter baumannii
☐ Bacteroides fragilis
☐ Citrobacter braakii/freundii
☐ Citrobacter koseri
☐ Enterobacter cloacae
☐ Enterococcus spp.
☐ Escherichia coli
☐ Klebsiella aerogenes
☐ K. oxytoca/michiganensis
☐ Klebsiella pneumoniae
☐ Morganella morganii
☐ Proteus mirabilis
☐ Pseudomonas aeruginosa
☐ Staphylococcus aureus
☐ Staphylococcus epidermidis
☐ Staphylococcus saprophyticus
☐ Strep. pyogenes (Group A)
☐ Varicella Zoster (Shingles)

ABX Resistance Markers

☐ β -lactamase (blaKPC)
☐ β -lactamase (CTX-M-Group 1)
☐ metallo- β -lactamase (blaNDM)
☐ Fluoroquinolones
☐ Methicillin/Oxacillin (mecA)
☐ Sulfonamides
☐ Trimethoprim

ICD 10 CODES

☐ L08.9 Local infection of the skin and subcutaneous tissue, unspecified
☐ Z22.39 Carrier of other specified bacterial disease
☐ Z22.322 Carrier or suspected carrier of MRSA

SPECIMEN SOURCE

☐ Aspiration
☐ Other: _____☐ Culture ID w/ Reflexive Antimicrobial Susceptibility Testing (AST is not available for STI or Vaginitis)

ICD 10 CODES

☐ _____
☐ _____
☐ _____

STEP 6

☐ Vaginitis☐ Atopobium vaginae
☐ Bacteroides fragilis
☐ BVAB-2
☐ Candida albicans
☐ Candida dubliniensis
☐ Candida glabrata
☐ Candida krusei

ICD 10 CODES

☐ N76.0 Acute vaginitis
☐ N77.1 Vaginitis, vulvitis, & vulvovaginitis
☐ B37.3 Candidiasis of vulva & vagina
☐ Z30.9 Encounter for contraceptive management

SPECIMEN SOURCE

☐ Vaginal Swab

ICD 10 CODES

☐ N76.0 Acute vaginitis
☐ N77.1 Vaginitis, vulvitis, & vulvovaginitis
☐ B37.3 Candidiasis of vulva & vagina
☐ Z30.9 Encounter for contraceptive management

SPECIMEN SOURCE

☐ Vaginal Swab

ICD 10 CODES

☐ R19.7 Diarrhea, unspecified
☐ A06.0 Acute Dysentery
☐ R50.9 Fever
☐ E86.0 Dehydration

SPECIMEN SOURCE

☐ Rectal Swab
☐ Stool Specimen
☐ Other: _____

ICD 10 CODES

☐ Z22.322 Carrier or suspected carrier of MRSA
☐ Z16.19 Resistance to other specified Beta Lactam antibiotics☐ STI☐ Atopobium vaginae
☐ Chlamydia trachomatis
☐ Gardnerella vaginalis
☐ Haemophilus ducreyi
☐ HHV-1 & 2 (Herpes Simplex)
☐ Neisseria gonorrhoeae
☐ Treponema pallidum
☐ Trichomonas vaginalis

ICD 10 CODES

☐ N76.0 Acute vaginitis
☐ N89.8 Other specified noninflammatory disorders of vagina
☐ R36.9 Urinary discharge unspecified
☐ Z30.9 Encounter for contraceptive management

SPECIMEN SOURCE

☐ Urine
☐ Other: _____

ICD 10 CODES

☐ Adenovirus
☐ Astrovirus
☐ Campylobacter coli/jejuni/lari
☐ Clostridium difficile
☐ E. coli (VTEC, O157)
☐ Norovirus (GI/GII)
☐ Rotavirus
☐ Salmonella spp.
☐ Sapovirus
☐ Shigella spp.
☐ Yersinia enterocolitica

ICD 10 CODES

☐ R19.7 Diarrhea, unspecified
☐ A06.0 Acute Dysentery
☐ R50.9 Fever
☐ E86.0 Dehydration

SPECIMEN SOURCE

☐ Rectal Swab
☐ Stool Specimen
☐ Other: _____

ICD 10 CODES

☐ Z22.322 Carrier or suspected carrier of MRSA
☐ Z16.19 Resistance to other specified Beta Lactam antibiotics☐ Fungal Infection☐ Alternaria spp.
☐ Aspergillus spp.
☐ Fusarium spp.
☐ Scytalidium dimidiatum
☐ Sarcocladium strictum
☐ Candida albicans
☐ Candida glabrata
☐ Candida krusei
☐ Candida parapsilosis
☐ Candida tropicalis
☐ Cryptococcus spp.
☐ Malassezia spp.
☐ Meyerozyma guilliermondii
☐ Trichophyton anthropophilic spp.
☐ Trichophyton zoophilic spp.
☐ Microsporum canis

ICD 10 CODES

☐ B35.1 Onychomycosis
☐ _____
☐ _____

SPECIMEN SOURCE

☐ Nail Clipping
☐ Other: _____
☐ Skin Scraping

ICD 10 CODES

☐ B35.1 Onychomycosis
☐ _____
☐ _____

SPECIMEN SOURCE

☐ Nail Clipping
☐ Other: _____
☐ Skin Scraping

ICD 10 CODES

☐ B35.1 Onychomycosis
☐ _____
☐ _____

SPECIMEN SOURCE

☐ Nail Clipping
☐ Other: _____
☐ Skin Scraping

ICD 10 CODES

☐ B35.1 Onychomycosis
☐ _____
☐ _____

SPECIMEN SOURCE

☐ Nail Clipping
☐ Other: _____
☐ Skin Scraping

ICD 10 CODES

☐ B35.1 Onychomycosis
☐ _____
☐ _____

SPECIMEN SOURCE

☐ Nail Clipping
☐ Other: _____
☐ Skin Scraping

6 Please indicate if your patient has taken antibiotics in the past 72 hours.

☐ Yes ☐ NoFOR MOST UP TO DATE REQUISITION FORM, PLEASE VISIT WWW.STREAMLINESCI.COM/FORMS-SUMMARY-LINKS

SPECIMEN PACKAGING, TRANSPORT AND RESULT DELIVERY




SPECIMEN PACKAGING AND TRANSPORT

1. Ensure sample lids are secure to avoid leakage or contamination.
2. Place specimen into a provided biohazard bag.
3. Insert requisition form, including all patient insurance information and demographics, into the specimen biohazard bag's side pocket.
4. Place the biohazard bag containing the patient specimen into the provided lock box.
5. Enter a pickup request ticket through the online portal or contact Streamline Scientific customer service team to schedule a pickup (unless a nightly pickup schedule has been established).

RESULT DELIVERY

Results will be available via our online portal system. This portal can be accessed by visiting our website

Result delivery via Secure Fax or Secure email is available. If you would prefer either of these methods, please contact customer service for setup.



RESPIRATORY PANEL TARGET CATALOG

VIRAL PATHOGENS

- Adenovirus
- Bocavirus
- COVID-19 (SARS-CoV-2)
- Coronavirus 229E
- Coronavirus HKU1
- Coronavirus NL63
- Coronavirus OC43
- EBV (mononucleosis)
- Enterovirus
- Human Metapneumovirus A
- Human Metapneumovirus B
- Influenza A
- Influenza B
- Parainfluenza Virus type 1
- Parainfluenza Virus type 2
- Parainfluenza Virus type 3
- Parainfluenza Virus type 4
- Respiratory Syncytial Virus
- Rhinovirus

BACTERIAL PATHOGENS

- Acinetobacter baumannii*
- Bordetella pertussis
- Chlamydia pneumoniae
- Enterobacter cloacae*
- Haemophilus influenzae*
- Klebsiella aerogenes*
- Klebsiella pneumoniae*
- Legionella pneumophila
- Moraxella catarrhalis*
- Mycoplasma pneumoniae
- Proteus mirabilis*
- Pseudomonas aeruginosa*
- Staphylococcus aureus*
- Staphylococcus epidermidis
- Streptococcus pneumoniae*
- Streptococcus pyogenes (Group A)**

ABX RESISTANCE MARKER

- Methicillin/Oxacillin (mecA)

*Limitation: Test does not differentiate between a patient with acute infection or an asymptomatic carrier.

**S. pyogenes detected from a throat swab is diagnostic of pharyngitis; S. pyogenes detected from a nasopharyngeal swab could indicate an asymptomatic carrier.

For panel offerings, please visit streamlinesci.com

RESPIRATORY INFECTION SAMPLE COLLECTION

Follow the instructions below depending on the specimen source:




NASOPHARYNGEAL SWAB:

1. Tilt the patient's head back 70 degrees.
2. Insert the swab into the nostril (the swab should reach depth equal to the distance from nostril to outer opening of the ear). Leave the swab in place for several seconds to absorb secretions.
3. Slowly remove the swab while rotating it.



MID-TURBINATE (PEDIATRIC) SWAB:

1. Gently insert the swab into the nostril.
2. Using a gentle rotation, push the swab until a slight resistance is met at the level of turbinates.
3. Rotate the swab several times against the nasal wall.



OROPHARYNGEAL SWAB:

1. Insert the swab into the posterior pharynx and tonsillar areas.
2. Rub the swab over both tonsillar pillars and posterior oropharyngeal and avoid touching the tongue, teeth, and gums.



SPUTUM:

1. Educate the patient about the difference between sputum and oral secretions.
2. Have the patient rinse the mouth with water and then expectorate deep cough sputum directly into a sterile screw-cap collection cup or sterile dry container.
3. Transfer this sputum to a tube with a liquid buffer for transport.

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310142
Date Collected: 11/08/23 3:09:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Controls

Panel Positive Control ¹ **PASS**
Panel Negative Control ² **PASS**

(1) Positive control is synthetic inactive pathogen
(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template
(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Test Results	Comments
COVID-19 Collection Type: Nasopharyngeal Swab		
COVID-19	Not Detected	[11/08/23] COVID-19 assay reviewed and approved under FDA Emergency Use Authorization #200522.
CT Value	0	[11/08/23] CT value indicates the number of amplification cycles by real-time PCR needed to detect specific sequences in SARS-CoV-2. CT values are inversely proportional to the amount of target nucleic acid in the sample (i.e. the lower the CT level, the greater the amount of target nucleic acid in the sample). CT values of 0 indicate no SARS-CoV-2 detected. The reference range for Streamline Scientific RT-PCR assays is <40 cycles of amplification.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY COVID Respiratory Plus Collection Type: Nasopharyngeal Swab			
Bocavirus	DETECTED - LOW	< 1,000	[11/08/23] Assay is developed to detect all strains of this pathogen
Pseudomonas aeruginosa	DETECTED - MEDIUM	1,000 - 100,000	[11/08/23] Limitation: Test does not differentiate between patient with acute infection or an asymptomatic carrier.
Streptococcus pneumoniae	DETECTED - HIGH	> 100,000	[11/08/23] Limitation: Test does not differentiate between patient with acute infection or an asymptomatic carrier.

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE RESPIRATORY PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310142
Date Collected: 11/08/23 3:09:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
COVID Respiratory Plus		Collection Type: Nasopharyngeal Swab	
Acinetobacter baumannii	Not Detected	Not Detected	
Adenovirus	Not Detected	Not Detected	[11/08/23] Assay is developed to detect all strains of this pathogen
Bocavirus	DETECTED - LOW	< 1,000	[11/08/23] Assay is developed to detect all strains of this pathogen
Bordetella pertussis	Not Detected	Not Detected	[11/08/23] This test does not differentiate between B. pertussis and B. holmesii.
Chlamydomphila pneumoniae	Not Detected	Not Detected	
Coronavirus 229E	Not Detected	Not Detected	
Coronavirus HKU1	Not Detected	Not Detected	
Coronavirus NL63	Not Detected	Not Detected	
Coronavirus OC43	Not Detected	Not Detected	
EBV (Mononucleosis)	Not Detected	Not Detected	
Enterobacter cloacae	Not Detected	Not Detected	
Enterovirus	Not Detected	Not Detected	[11/08/23] Enterovirus includes Coxsackievirus types A9, A10, A16, B5, and Echovirus serotypes [11/08/23] May cross-react with Rhinovirus (types A and B)
Haemophilus influenzae	Not Detected	Not Detected	
HMPV A (Human Metapneumovirus)	Not Detected	Not Detected	
HMPV B (Human Metapneumovirus)	Not Detected	Not Detected	
Influenza A	Not Detected	Not Detected	
Influenza B	Not Detected	Not Detected	
Klebsiella aerogenes	Not Detected	Not Detected	
Klebsiella pneumoniae	Not Detected	Not Detected	
Legionella pneumophila	Not Detected	Not Detected	
mecA (Methicillin/Oxacillin resistance)	Not Detected	Not Detected	
Moraxella catarrhalis	Not Detected	Not Detected	
Mycoplasma pneumoniae	Not Detected	Not Detected	
Parainfluenza 1	Not Detected	Not Detected	
Parainfluenza 2	Not Detected	Not Detected	
Parainfluenza 3	Not Detected	Not Detected	
Parainfluenza 4	Not Detected	Not Detected	
Proteus mirabilis	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.

Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.

Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.

In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE RESPIRATORY PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310142
Date Collected: 11/08/23 3:09:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Pseudomonas aeruginosa	DETECTED - MEDIUM	1,000 - 100,000	[11/08/23] Limitation: Test does not differentiate between patient with acute infection or an asymptomatic carrier.
Rhinovirus (types A & B)	Not Detected	Not Detected	[11/08/23] Assay is developed to detect all strains of this pathogen [11/08/23] May cross-react with Enterovirus
Respiratory Syncytial Virus	Not Detected	Not Detected	
Staphylococcus aureus	Not Detected	Not Detected	
Staphylococcus epidermidis	Not Detected	Not Detected	
Streptococcus pneumoniae	DETECTED - HIGH	> 100,000	[11/08/23] Limitation: Test does not differentiate between patient with acute infection or an asymptomatic carrier.
Streptococcus pyogenes (Group A)	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 3

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE RESPIRATORY PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

UTI PANEL TARGETS

PATHOGENS

- Acinetobacter baumannii
- Bacteroides fragilis
- Candida albicans
- Candida dubliniensis
- Candida glabrata
- Candida krusei
- Candida parapsilosis
- Candida tropicalis
- Citrobacter braakii/freundii
- Citrobacter koseri
- Enterobacter cloacae
- Enterococcus spp.
- Escherichia coli
- Klebsiella aerogenes
- K. oxytoca/michiganensis
- Klebsiella pneumoniae
- Morganella morganii
- Mycoplasma genitalium
- Mycoplasma hominis
- Prevotella bivia
- Proteus mirabilis
- Pseudomonas aeruginosa
- Serratia marcescens
- Staphylococcus aureus
- Staphylococcus epidermidis
- Staphylococcus saprophyticus
- Streptococcus agalactiae (Group B)
- Streptococcus pyogenes (Group A)
- Ureaplasma urealyticum

ABX RESISTANCE MARKER

- β -lactamase (blaKPC)
- β -lactamase (CTX-M-Group 1)
- Metallo- β -lactamase (blaNDM)
- Fluoroquinolones
- Methicillin/Oxacillin (mecA)
- Sulfonamides
- Trimethoprim

For panel offerings, please visit streamlinesci.com

UTI SAMPLE COLLECTION

Collect the patient sample using one of the procedures below:

URINE SAMPLE:

If submitting a urine sample: Collect a urine sample from the patient in a sterile urine collection cup with lid-integrated transfer device using the following gender-specific instructions:

Female Urine Sample

1. Patient should use a packaged, moist towel to clean the vulva and perianal area starting from front to back. Repeat with a second moist towel.
2. Patient should then spread their labia with one hand and start urinating into the toilet. Within the other hand, they should put the urine container under the genital area to catch the stream of urine without touching any skin.

Male Urine Sample

1. Patient should retract the foreskin from the penis if necessary and use the packaged towel to clean the penis from the tip to the base. Repeat with a second moist towel.
2. Patient should retract the foreskin if necessary with one hand and start urinating into the toilet. Then, position the urine container with the other hand to catch the stream without touching any skin.

For PCR testing with culture request

Replace and tighten the collection cup lid, and transfer the urine sample into a vacutainer tube with preservative:

1. Remove the sticker from the lid to access the integrated transfer device.
2. Insert the vacutainer tube vertically into the transfer device, puncturing the rubber septum and allowing the tube to fill with urine.
3. Remove the vacutainer tube and discard the collection cup. Mix urine and preservative by gently inverting the tube 8-10 times.

For PCR testing only

Urine Specimen Collection Instructions

Urine should be collected in the provided sterile cup. Next, use the provided disposable transfer pipette to transfer at least 1mL of urine into the tube. Replace the specimen lid, ensure it is properly labeled with the patient's name and DOB and send to the lab. Urine collection cup and transfer pipette can be disposed of in biohazard waste.



CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310077
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Controls

Panel Positive Control ¹ PASS
Panel Negative Control ² PASS

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY UTI Pathogens Plus			Collection Type: Urine clean catch
Candida krusei	DETECTED - LOW	< 1,000	
Escherichia coli	DETECTED - HIGH	> 100,000	
Serratia marcescens	DETECTED - MEDIUM	1,000 - 10,000	

Class (Gene Name)	Lab Result (Qualitative)	Resistance Gene Targets Identified	Associated Resistances (Antibiotics to Avoid)
SUMMARY UTI Antibiotic Resistance Markers			Collection Type: Urine clean catch
vanA (Vancomycin)	DETECTED	vanA	[11/09/23]Vancomycin

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE UTI PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310077
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
UTI Pathogens Plus			
Collection Type: Urine clean catch			
Acinetobacter baumannii	Not Detected	Not Detected	
Bacteroides fragilis	Not Detected	Not Detected	
Candida albicans	Not Detected	Not Detected	
Candida dubliniensis	Not Detected	Not Detected	
Candida glabrata	Not Detected	Not Detected	
Candida krusei	DETECTED - LOW	< 1,000	
Candida parapsilosis	Not Detected	Not Detected	
Candida tropicalis	Not Detected	Not Detected	
Citrobacter freundii/braakii	Not Detected	Not Detected	
Citrobacter koseri	Not Detected	Not Detected	
Enterobacter cloacae	Not Detected	Not Detected	
Enterococcus spp.	Not Detected	Not Detected	
Escherichia coli	DETECTED - HIGH	> 100,000	
Klebsiella aerogenes	Not Detected	Not Detected	
Klebsiella oxytoca/michiganensis	Not Detected	Not Detected	
Klebsiella pneumoniae	Not Detected	Not Detected	
Morganella morganii	Not Detected	Not Detected	
Mycoplasma genitalium	Not Detected	Not Detected	
Mycoplasma hominis	Not Detected	Not Detected	
Prevotella bivia	Not Detected	Not Detected	
Proteus mirabilis	Not Detected	Not Detected	
Pseudomonas aeruginosa	Not Detected	Not Detected	
Serratia marcescens	DETECTED - MEDIUM	1,000 - 10,000	
Staphylococcus aureus	Not Detected	Not Detected	
Staphylococcus epidermidis	Not Detected	Not Detected	
Staphylococcus saprophyticus	Not Detected	Not Detected	
Streptococcus agalactiae (GBS)	Not Detected	Not Detected	
Streptococcus pyogenes (Group A)	Not Detected	Not Detected	
Ureaplasma urealyticum	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE UTI PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310077
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Class (Gene Name)	Lab Result (Qualitative)	Resistance Gene Targets Identified	Associated Resistances (Antibiotics to Avoid)
UTI Antibiotic Resistance Markers		Collection Type: Urine clean catch	
Class A Beta-lactamase (blaKPC)	Not Detected	KPC-2-8,10,11,13-22,24-33	[11/09/23] Carbapenems, Cephalosporins, Penicillins, Beta-lactamase inhibitors, Aztreonam
Class A Beta-lactamase (CTX-M-Group 1)	Not Detected	blaCTX-M-1,3,10,12,15,22,23,28; blaFEC-1	[11/09/23] Cephalosporins, Penicillins, Aztreonam
Class B metallo Beta-lactamase (blaNDM)	Not Detected	NDM (1-21)	[11/09/23] Carbapenems, Cephalosporins, Penicillins, Beta-lactamase inhibitors
Fluoroquinolones	Not Detected	qnrS 1-5,7-9; qnrB Group 1; qnrB Group 5	[11/09/23] Ciprofloxacin, Gemifloxacin, Levofloxacin, Moxifloxacin, Norfloxacin, Ofloxacin
mecA (Methicillin/Oxacillin resistance)	Not Detected	mecA	[11/09/23] Oxacillin
Sulfonamides	Not Detected	sul1; sul2; sul3	[11/09/23] Sulfadiazine, Sulfamethizole, Sulfamethoxazole, Sulfasalazine, Sulfisoxazole
Trimethoprim	Not Detected	dfrA1; dfrA5; dfrA11; dfrA17	[11/09/23] Primsol
vanA (Vancomycin)	DETECTED	vanA	[11/09/23] Vancomycin
vanB (Vancomycin)	Not Detected	vanB	[11/09/23] Vancomycin

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 3

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.

Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.

Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.

In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE UTI PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

WOUND PANEL

PATHOGENS

- Acinetobacter baumannii
- Bacteroides fragilis
- Citrobacter braakii/freundii
- Citrobacter koseri
- Enterobacter cloacae
- Enterococcus spp.
- Escherichia coli
- Klebsiella aerogenes
- K. oxytoca/michiganensis
- Klebsiella pneumoniae
- Morganella morganii
- Proteus mirabilis
- Pseudomonas aeruginosa
- Staphylococcus aureus
- Staphylococcus epidermidis
- Staphylococcus saprophyticus
- Streptococcus pyogenes (Group A)
- Varicella Zoster (Shingles)

ABX RESISTANCE MARKER

- β -lactamase (blaKPC)
- β -lactamase (CTX-M-Group 1)
- Metallo- β -lactamase (blaNDM)
- Fluoroquinolones
- Methicillin/Oxacillin (mecA)
- Sulfonamides
- Trimethoprim

For panel offerings, please visit streamlinesci.com

WOUND/INFECTION SAMPLE COLLECTION

Collect the patient sample using one of the procedures below:

ASPIRATION

1. The surface of the wound/abscess should be carefully cleansed and debrided using sterile gauze and saline before attempting to aspirate the specimen.
2. Aspirate the specimen and place 0.5 to 1.0 mL of the aspirate directly into a transport tube.
3. Without contaminating the included swab, place the swab into the transport tube all the way to the bottom. Break the swab at the scored breakpoint indication line and leave the bottom portion inside the transport tube, partially submerged in buffer solution. Screw the top onto the tube tightly to secure the specimen.

SWAB

1. Cleanse and debride the wound with sterile gauze and saline.
2. Exudate and brushings of the wound base (including advancing margins) should be collected using the swab.
3. Without contaminating the swab, place the swab into the transport tube all the way to the bottom.
4. Break the swab at the scored breakpoint indication line and leave the bottom portion inside the transport tube, partially submerged in buffer solution. Screw the top onto the tube tightly to secure the specimen.

Note: If culture ID is requested, sample should be sent in Liquid Amies media.



CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310076
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Controls

Panel Positive Control ¹ **PASS**
Panel Negative Control ² **PASS**

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY Wound/Dermatology Panel			Collection Type: Lesion
Escherichia coli	DETECTED - HIGH	> 100,000	
Staphylococcus epidermidis	DETECTED - LOW	< 1,000	[11/09/23] Low detection of S. epidermidis may be indicative of commensal flora.
Varicella Zoster virus	DETECTED	DETECTED	

Class (Gene Name)	Lab Result (Qualitative)	Resistance Gene Targets Identified	Associated Resistances (Antibiotics to Avoid)
SUMMARY Wound Antibiotic Resistance Markers			Collection Type: Lesion
mecA (Methicillin/Oxacillin resistance)	DETECTED	mecA	[11/09/23]Oxacillin

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE WOUND/INFECTION PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310076
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Wound/Dermatology Panel		Collection Type: Lesion	
Acinetobacter baumannii	Not Detected	Not Detected	
Bacteroides fragilis	Not Detected	Not Detected	
Citrobacter freundii/braakii	Not Detected	Not Detected	
Citrobacter koseri	Not Detected	Not Detected	
Enterobacter cloacae	Not Detected	Not Detected	
Enterococcus spp.	Not Detected	Not Detected	
Escherichia coli	DETECTED - HIGH	> 100,000	
Klebsiella aerogenes	Not Detected	Not Detected	
Klebsiella oxytoca/michiganesis	Not Detected	Not Detected	
Klebsiella pneumoniae	Not Detected	Not Detected	
Morganella morganii	Not Detected	Not Detected	
Proteus mirabilis	Not Detected	Not Detected	
Pseudomonas aeruginosa	Not Detected	Not Detected	
Staphylococcus aureus	Not Detected	Not Detected	
Staphylococcus epidermidis	DETECTED - LOW	< 1,000	[11/09/23] Low detection of S. epidermidis may be indicative of commensal flora.
Staphylococcus saprophyticus	Not Detected	Not Detected	
Streptococcus pyogenes (Group A)	Not Detected	Not Detected	
Varicella Zoster virus	DETECTED	DETECTED	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE WOUND/INFECTION PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11092310076
Date Collected: 11/09/23 9:17:00 AM
Date Accessioned: 11/09/2023
Date Reported: 11/09/2023
Faxed to: 18777966185

Class (Gene Name)	Lab Result (Qualitative)	Resistance Gene Targets Identified	Associated Resistances (Antibiotics to Avoid)
Wound Antibiotic Resistance Markers		Collection Type: Lesion	
Class A Beta-lactamase (blaKPC)	Not Detected	KPC-2-8,10,11,13-22,24-33	[11/09/23] Carbapenems, Cephalosporins, Penicillins, Beta-lactamase inhibitors, Aztreonam
Class A Beta-lactamase (CTX-M-Group 1)	Not Detected	blaCTX-M-1,3,10,12,15,22,23,28; blaFEC-1	[11/09/23] Cephalosporins, Penicillins, Aztreonam
Class B metallo Beta-lactamase (blaNDM)	Not Detected	NDM (1-21)	[11/09/23] Carbapenems, Cephalosporins, Penicillins, Beta-lactamase inhibitors
Fluoroquinolones	Not Detected	qnrS 1-5,7-9; qnrB Group 1; qnrB Group 5	[11/09/23] Ciprofloxacin, Gemifloxacin, Levofloxacin, Moxifloxacin, Norfloxacin, Ofloxacin
mecA (Methicillin/Oxacillin resistance)	DETECTED	mecA	[11/09/23] Oxacillin
Sulfonamides	Not Detected	sul1; sul2; sul3	[11/09/23] Sulfadiazine, Sulfamethizole, Sulfamethoxazole, Sulfasalazine, Sulfisoxazole
Trimethoprim	Not Detected	dfrA1; dfrA5; dfrA11; dfrA17	[11/09/23] Primsol
vanA (Vancomycin)	Not Detected	vanA	[11/09/23] Vancomycin
vanB (Vancomycin)	Not Detected	vanB	[11/09/23] Vancomycin

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 3

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE WOUND/INFECTION PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

STI PANEL

PATHOGENS

- Atopobium vaginae
- Chlamydia trachomatis
- Gardnerella vaginalis
- Haemophilus ducreyi
- HHV-1 (Herpes Simplex)
- HHV-2 (Herpes Simplex)
- Neisseria gonorrhoeae
- Treponema pallidum
- Trichomonas vaginalis

For panel offerings, please visit streamlinesci.com

STI SAMPLE COLLECTION

If submitting a urine sample: Collect a urine sample from the patient in a sterile urine collection cup and transfer to a vacutainer tube with bacteriostatic preservative.

Note: Clean-catch urine is not recommended for STI or Candida testing.

If submitting a swab sample: Collect a swab sample from the patient with a synthetic flocked swab by following the instructions below depending on the specimen source:

Vaginal swab:

1. Remove the swab applicator and collect a specimen by rotating the swab against the wall of the vaginal canal several times for 20-30 seconds.
2. Withdraw the swab without touching the vaginal surface.

Urethral swab:

1. Gently insert the swab into the urethra (1-2 cm for women, 2-4 cm for men).
2. Rotate the swab in one direction for a minimum of 10 seconds.
3. Withdraw the swab.

*If an ulcer is the intended specimen source, please refer to wound specimen collection instructions

Valid specimen type for Chlamydia trachomatis/Neisseria gonorrhoeae/Trichomonas vaginalis testing only:

Throat swab (Oropharyngeal swab):

1. Insert the swab into the posterior pharynx and tonsillar areas.
2. Rub the swab over both tonsillar pillars and posterior oropharyngeal. Avoid touching the tongue, teeth, and gums.

Rectal swab:

1. Insert a sterile swab approximately 2.5 cm into the anal canal.
2. Move the swab from side to side in the anal canal. Allow the swab to remain 10-30 seconds for the absorption of organisms onto the swab.
3. Remove the swab and insert it into a vial containing 1-3ml of transport media.

Disclaimer/Collection - Note:

PCR tests can be used to confirm a suspicion of the presence of sexually transmitted infections in the routine clinical setting. However, in cases of rape or sexual abuse in children under the age of 15, confirmatory testing (along with the appropriate chain of custody as outlined by the CDC) should be performed as recommended by the CDC.

When collecting a urethral swab or urine specimen from a male or female patient, the patient should not have urinated for **at least an hour** before the specimen is collected. The first-morning urine specimen is preferred due to a large number of cells usually present.

NOTE: Certain organisms are intracellular; therefore, there must be enough human cells present to detect the organism.



CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310192
Date Collected: 11/08/23 10:41:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Controls

Panel Positive Control ¹ **PASS**
Panel Negative Control ² **PASS**

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY Sexually Transmitted Infection Pathogens		Collection Type: Urine	
<i>Gardnerella vaginalis</i>	DETECTED - LOW	< 1,000	
<i>Neisseria gonorrhoeae</i>	DETECTED	DETECTED	[11/08/23] N.gonorrhoeae was detected by PCR and is a Notifiable Pathogen per ADPH. Result to be confirmed by lab before reporting to ADPH. The required time for the ordering provider to notify ADPH is within 5 days from receipt of report.

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.

Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.

Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.

In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE STI PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207
BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST
BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310192
Date Collected: 11/08/23 10:41:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Sexually Transmitted Infection Pathogens		Collection Type: Urine	
Atopobium vaginae	Not Detected	Not Detected	
Chlamydia trachomatis	Not Detected	Not Detected	
Gardnerella vaginalis	DETECTED - LOW	< 1,000	
Haemophilus ducreyi	Not Detected	Not Detected	
HHV-1 (Herpes Simplex Virus)	Not Detected	Not Detected	
HHV-2 (Herpes Simplex Virus)	Not Detected	Not Detected	
Neisseria gonorrhoeae	DETECTED	DETECTED	[11/08/23] N.gonorrhoeae was detected by PCR and is a Notifiable Pathogen per ADPH. Result to be confirmed by lab before reporting to ADPH. The required time for the ordering provider to notify ADPH is within 5 days from receipt of report.
Treponema pallidum	Not Detected	Not Detected	
Trichomonas vaginalis	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE STI PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

VAGINITIS PANEL

PATHOGENS

- Atopobium vaginae
- Bacteroides fragilis
- BVAB-2
- Candida albicans
- Candida dubliniensis
- Candida glabrata
- Candida krusei
- Candida lusitanae
- Candida parapsilosis
- Candida tropicalis
- Chlamydia trachomatis
- Enterococcus spp.
- Escherichia coli
- Gardnerella vaginalis
- Haemophilus ducreyi
- HHV-1 (Herpes Simplex)
- HHV-2 (Herpes Simplex)
- Lactobacillus crispatus
- Lactobacillus gasseri
- Lactobacillus iners
- Lactobacillus jensenii
- Megasphaera Type 1
- Megasphaera Type 2
- Mobiluncus curtisii
- Mobiluncus mulieris
- Mycoplasma genitalium
- Mycoplasma hominis
- Neisseria gonorrhoeae
- Prevotella bivia
- Staphylococcus aureus
- Streptococcus agalactiae (Group B)
- Treponema pallidum
- Trichomonas vaginalis
- Ureaplasma urealyticum

VAGINITIS SAMPLE COLLECTION



Collect a vaginal sample from the patient using a synthetic flocked swab by inserting the swab into the vagina and turn the swab 3 times.

.....

Disclaimer/Collection Note:

PCR tests can be used to confirm a suspicion of the presence of sexually transmitted infections in the routine clinical setting. However, in cases of rape or sexual abuse in children under the age of 15, confirmatory testing (along with the appropriate chain of custody as outlined by the CDC) should be performed as recommended by the CDC.

NOTE: Certain organisms are intracellular; therefore, there must be enough human cells present to detect the organism.

CLINIC INFORMATION

 Name: Streamline Pathology Laboratory
 Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
 Provider: Test, Doctor MD

PATIENT INFORMATION

 Name: TEST, PATIENT
 DOB: 3/31/1995 Sex: M
 Phone: (777)777-7777
 Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

 Lab Accession Number: S-11082310190
 Date Collected: 11/08/23 10:24:00 PM
 Date Accessioned: 11/08/2023
 Date Reported: 11/08/2023
 Faxed to: 18777966185

Controls

 Panel Positive Control ¹ **PASS**
 Panel Negative Control ² **PASS**

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY Vaginitis Pathogens		Collection Type: Vaginal swab (specimen)	
BVAB-2	DETECTED - LOW	< 1,000	
Candida parapsilosis	DETECTED - MEDIUM	1,000 - 10,000	
Lactobacillus jensenii	DETECTED - MEDIUM	10,000 - 100,000	
Prevotella bivia	DETECTED - HIGH	> 100,000	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
 Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
 Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
 In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

 FOR MORE VAGINITIS PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310190
Date Collected: 11/08/23 10:24:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Vaginitis Pathogens			
Collection Type: Vaginal swab (specimen)			
Atopobium vaginae	Not Detected	Not Detected	
Bacteroides fragilis	Not Detected	Not Detected	
BVAB-2	DETECTED - LOW	< 1,000	
Candida albicans	Not Detected	Not Detected	
Candida dubliniensis	Not Detected	Not Detected	
Candida glabrata	Not Detected	Not Detected	
Candida krusei	Not Detected	Not Detected	
Candida lusitanae	Not Detected	Not Detected	
Candida parapsilosis	DETECTED - MEDIUM	1,000 - 10,000	
Candida tropicalis	Not Detected	Not Detected	
Chlamydia trachomatis	Not Detected	Not Detected	
Enterococcus spp.	Not Detected	Not Detected	
Escherichia coli	Not Detected	Not Detected	
Gardnerella vaginalis	Not Detected	Not Detected	
Haemophilus ducreyi	Not Detected	Not Detected	
HHV-1 (Herpes Simplex Virus)	Not Detected	Not Detected	
HHV-2 (Herpes Simplex Virus)	Not Detected	Not Detected	
Lactobacillus crispatus	Not Detected	Not Detected	
Lactobacillus gasseri	Not Detected	Not Detected	
Lactobacillus iners	Not Detected	Not Detected	
Lactobacillus jensenii	DETECTED - MEDIUM	10,000 - 100,000	
Megasphaera Type 1	Not Detected	Not Detected	
Megasphaera Type 2	Not Detected	Not Detected	
Mobiluncus curtisii	Not Detected	Not Detected	
Mobiluncus mulieris	Not Detected	Not Detected	
Mycoplasma genitalium	Not Detected	Not Detected	
Mycoplasma hominis	Not Detected	Not Detected	
Neisseria gonorrhoeae	Not Detected	Not Detected	
Prevotella bivia	DETECTED - HIGH	> 100,000	
Staphylococcus aureus	Not Detected	Not Detected	
Streptococcus agalactiae (GBS)	Not Detected	Not Detected	
Treponema pallidum	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE VAGINITIS PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology
Laboratory
Address: 2868 Acton Road
Suite 207
BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST
BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310190
Date Collected: 11/08/23 10:24:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Trichomonas vaginalis	Not Detected	Not Detected	
Ureaplasma urealyticum	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 3

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE VAGINITIS PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

GASTROINTESTINAL PANEL

PATHOGENS

Viral Pathogens:

- Adenovirus F40/F41
- Astrovirus
- Norovirus GI/GII
- Rotavirus A

Bacterial Pathogens:

- *C. difficile* (tcdA, tcdB)
- *Campylobacter* (*C. coli*, *C. jejuni*, and *C. lari*)
- Enteroinvasive *E. coli* (EIEC)/*Shigella*
- Enterotoxigenic *E. coli* (ETEC)
- *Salmonella* spp.
- Shiga-like toxin-producing *E. coli* (STEC)
- *Vibrio* spp.
- *Yersinia enterocolitica*

Parasites:

- *Cryptosporidium* spp.
- *Entamoeba histolytica*
- *Giardia lamblia*

For panel offerings, please visit streamlinesci.com

GASTROINTESTINAL SAMPLE COLLECTION

Collect a gastrointestinal stool specimen sample or a rectal swab from the patient using a swab with Cary Blair buffer. Use the Cary Blair buffer specimen collection swab for the transport of either specimen swab or rectal swab by using the following instructions:

1. Remove the swab and transport tube from the packaging. Do not contaminate.
2. For stool specimen swabs, carefully collect a portion of the specimen by placing the swab tip directly into the specimen.
3. For rectal swabs, insert the tip of the swab approximately 1 inch beyond the anal sphincter.
4. Carefully rotate the swab to sample the anal crypts, then withdraw the swab.

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310130
Date Collected: 11/08/23 2:26:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Controls

Panel Positive Control ¹ PASS
Panel Negative Control ² PASS

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY Gastrointestinal Panel w/ C.Diff Add-On			Collection Type: Stool - Fecal
Clostridium difficile (tcdA/tcdB)	DETECTED - LOW	< 1,000	[11/08/23] * Due to the high prevalence of asymptomatic carriage of toxigenic C. difficile in infants, testing for CDI should never be routinely recommended for neonates or infants 12 months of age or younger with diarrhea * C. difficile testing should not be routinely performed in children with diarrhea who are 1-2 years of age unless other infectious or noninfectious causes have been excluded * In children 2 years of age or older, C. difficile testing is recommended for patients with prolonged or worsening diarrhea and risk factors (eg, underlying inflammatory bowel disease or immunocompromising conditions) or relevant exposures (eg, contact with the healthcare system or recent antibiotics)
Norovirus GI/GII	DETECTED	DETECTED	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.

Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.

Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.

In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE GASTROINTESTINAL PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310130
Date Collected: 11/08/23 2:26:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Gastrointestinal Panel w/ C.Diff Add-On		Collection Type: Stool - Fecal	
Adenovirus F40/F41	Not Detected	Not Detected	
Astrovirus	Not Detected	Not Detected	
Campylobacter coli/jejuni/lari/upsaliensis	Not Detected	Not Detected	
Clostridium difficile (tcdA/tcdB)	DETECTED - LOW	< 1,000	<p>[11/08/23] * Due to the high prevalence of asymptomatic carriage of toxigenic C. difficile in infants, testing for CDI should never be routinely recommended for neonates or infants 12 months of age or younger with diarrhea</p> <p>* C. difficile testing should not be routinely performed in children with diarrhea who are 1-2 years of age unless other infectious or noninfectious causes have been excluded</p> <p>* In children 2 years of age or older, C. difficile testing is recommended for patients with prolonged or worsening diarrhea and risk factors (eg, underlying inflammatory bowel disease or immunocompromising conditions) or relevant exposures (eg, contact with the healthcare system or recent antibiotics)</p>
Cryptosporidium spp.	Not Detected	Not Detected	
Entamoeba histolytica	Not Detected	Not Detected	
Enterotoxigenic E.coli (ETEC)	Not Detected	Not Detected	
Enteroinvasive E.coli (EIEC)/Shigella spp.	Not Detected	Not Detected	
Shiga-like Toxin producing E.coli (STEC)	Not Detected	Not Detected	
Giardia lamblia	Not Detected	Not Detected	
Norovirus GI/GII	DETECTED	DETECTED	
Rotavirus A	Not Detected	Not Detected	
Salmonella spp.	Not Detected	Not Detected	
Vibrio spp.	Not Detected	Not Detected	
Yersinia enterocolitica	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE GASTROINTESTINAL PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

FUNGAL INFECTION PANEL

PATHOGENS

- *Alternaria* spp.
- *Aspergillus* spp.
- *Fusarium* spp.
- *Scytalidium dimidiatum*
- *Sarocladium strictum*
- *Candida albicans*
- *Candida glabrata*
- *Candida krusei*
- *Candida parapsilosis*
- *Candida tropicalis*
- *Cryptococcus* spp.
- *Malassezia* spp.
- *Meyerozyma guilliermondii*
- *Trichophyton anthropophilic* spp.
- *Trichophyton zoophilic* spp.
- *Mircosporum canis*

BACTERIA ADD ON

- *Pseudomonas aeruginosa*

ABX RESISTANCE MARKER

- Methicillin/Oxacillin (*mecA*)

FUNGAL SPECIMEN COLLECTION

ACCEPTABLE SPECIMENS:

Nail clippings and skin scrapings in a dry collection tube. If a wound is infected, the area should be collected by a swab in liquid amies media.



DIRECTIONS FOR COLLECTING A NAIL CLIPPING

1. Wipe the nail collection site with 70% isopropyl alcohol.
2. Debride and discard nail clippings.
3. Obtain specimen from the most proximal area of nail and hyponychium. Minimum specimen amount size of nail and subungual debris is 3mm to 6mm (small pieces to obtain this size are preferred).
4. Use a curette to obtain any additional subungual debris.
5. Place the dry nail sample and debris into a dry sterile tube.

DIRECTIONS FOR A SKIN SCRAPING

1. Remove any traces of skin products, medications, or surface contaminants by wiping the area with a 70% isopropyl alcohol wipe.
2. Choose the best area to scrape by determining where fungal growth is most active.
3. Scrape the skin using a scalpel held at a blunt angle into dry collection tube. The greater amount of specimen, the better the result.

DIRECTIONS FOR A WOUND COLLECTION

1. Cleanse and debride the infected area with 70% isopropyl alcohol using sterile gauze. Saline can be used if there is an open wound.
2. Exudate and brushing of the base of the infected area (including advancing margins) should be collected using the swab. Without contaminating the swab, place the swab in the media.

CLINIC INFORMATION

 Name: Streamline Pathology Laboratory
 Address: 2868 Acton Road
 Suite 207
 BIRMINGHAM, AL 35243
 Provider: Test, Doctor MD

PATIENT INFORMATION

 Name: TEST, PATIENT
 DOB: 3/31/1995 Sex: M
 Phone: (777)777-7777
 Address: 456 MAPLE ST
 BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

 Lab Accession Number: S-11082310182
 Date Collected: 11/08/23 9:52:00 PM
 Date Accessioned: 11/08/2023
 Date Reported: 11/08/2023
 Faxed to: 18777966185

Controls

 Panel Positive Control ¹ **PASS**
 Panel Negative Control ² **PASS**

(1) Positive control is synthetic inactive pathogen

(2) Negative Control contains primers, probe, and enzymes with no DNA/RNA template

(3) A "Detected" result indicates the presence of a pathogen (99.99% confidence) above the assay cutoff.

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
SUMMARY Fungal Infection Panel w/ Bacterial Add-On_v2		Collection Type: Nail specimen (specimen)	
Cryptococcus spp.	DETECTED - MEDIUM	1,000 - 100,000	
Microsporum canis	DETECTED - HIGH	> 100,000	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
 Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
 Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
 In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

 FOR MORE FUNGAL INFECTION PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CLINIC INFORMATION

Name: Streamline Pathology Laboratory
Address: 2868 Acton Road
Suite 207
BIRMINGHAM, AL 35243
Provider: Test, Doctor MD

PATIENT INFORMATION

Name: TEST, PATIENT
DOB: 3/31/1995 Sex: M
Phone: (777)777-7777
Address: 456 MAPLE ST
BIRMINGHAM, AL 35243

SPECIMEN INFORMATION

Lab Accession Number: S-11082310182
Date Collected: 11/08/23 9:52:00 PM
Date Accessioned: 11/08/2023
Date Reported: 11/08/2023
Faxed to: 18777966185

Test Performed	Lab Result (3) (Qualitative Low/Medium/High)	DNA Copy Number	Comments
Fungal Infection Panel w/ Bacterial Add-On_v2		Collection Type: Nail specimen (specimen)	
Alternaria spp.	Not Detected	Not Detected	
Aspergillus spp.	Not Detected	Not Detected	
Candida albicans	Not Detected	Not Detected	
Candida glabrata	Not Detected	Not Detected	
Candida krusei	Not Detected	Not Detected	
Candida parapsilosis	Not Detected	Not Detected	
Candida tropicalis	Not Detected	Not Detected	
Cryptococcus spp.	DETECTED - MEDIUM	1,000 - 100,000	
Curvularia spp.	Not Detected	Not Detected	
Epidermophyton floccosum	Not Detected	Not Detected	
Fusarium spp.	Not Detected	Not Detected	
Malassezia spp.	Not Detected	Not Detected	
Meyerozyma guilliermondii	Not Detected	Not Detected	
Microsporum canis	DETECTED - HIGH	> 100,000	
Sarocladium strictum	Not Detected	Not Detected	
Scytalidium dimidiatum	Not Detected	Not Detected	
Trichophyton anthropophilic spp.	Not Detected	Not Detected	
Trichophyton zoophilic spp.	Not Detected	Not Detected	
Trichosporon spp.	Not Detected	Not Detected	
Pseudomonas aeruginosa	Not Detected	Not Detected	
mecA (Methicillin/Oxacillin resistance)	Not Detected	Not Detected	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, PATIENT

Lab Director: Robert Thomas, MD

Ver. PP231013.1 Page: 2

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.

Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.

Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.

In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

FOR MORE FUNGAL INFECTION PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

CULTURE ID AND SENSITIVITY TESTING



Liquid Amies buffer tubes or bacteriostatic preservative urine tubes should be used for samples with culture ID requests.

.....

Culture ID and Sensitivity Testing must be performed within 48 hours of specimen collection.

.....

Typical turnaround time is 48-72 hours. This timeframe may be extended in cases of mixed culture results.

.....

Culture ID and Sensitivity testing is not offered for STI or Vaginitis specimens.

CLINIC INFORMATION

 Name: Streamline Pathology Laboratory
 Address: 2868 Acton Road Suite 207 BIRMINGHAM, AL 35243
 Provider: Test, Doctor MD

PATIENT INFORMATION

 Name: TEST, TEST
 DOB: 1/1/2000 Sex: M
 Phone: (405)000-0000
 Address: 123 Test St Apt 10 BIRMINGHAM, AL 35213

SPECIMEN INFORMATION

 Lab Accession Number: S-06062310028
 Date Collected: 06/05/23 2:49:00 PM
 Date Accessioned: 06/05/2023
 Date Reported: 06/06/2023
 Faxed to: 18777966185

Culture ID and Antibiotic Sensitivity

Summary			Collection Type: Urine clean catch
Antibiotic Sensitivity Testing	Minimum Inhibitory Concentration (ug/mL)	Susceptibility	Notes
Isolated Organism 1			
<i>Escherichia coli</i>			
Organism Semi-Quantification	> 100,000 cfu/mL ug/mL		
Trimethoprim-sulfamethoxazole	<=2 ug/mL	S	
Nitrofurantoin	<=32 ug/mL	S	
Ceftriaxone	<=0.5 ug/mL	S	
Ampicillin	<=8 ug/mL	S	
Ciprofloxacin	<=0.047 ug/mL	S	
Cefazolin	<=1 ug/mL	S	
Gentamicin	<=2 ug/mL	S	
Isolated Organism 2			
<i>Streptococcus agalactiae</i>			
Organism Semi-Quantification	10,000 cfu/mL-100,000 cfu/mL ug/mL		
Vancomycin	<=0.25 ug/mL	S	
Penicillin	<=0.12 ug/mL	S	
Clindamycin	>2 ug/mL	R	

Clinic Name: Streamline Pathology Laboratory

Patient name: TEST, TEST

Lab Director: Ty Thomas, MD

Page: 1

Test results must be evaluated with clinical symptoms to diagnose disease. All tests established and validated by Laboratory and not FDA approved unless otherwise indicated.
 Methodology statement: Real-time PCR assays are designed to detect pathogens with clinical significance and analytical sensitivity and specificity greater than 99%.
 Limitations: While these assays are very sensitive and specific, theoretically these assays could detect pathogens not listed, resulting in a false positive.
 In addition, while these assays are very specific, there may be target pathogen sequences with unknown sequence variability which may not be detected, resulting in a false negative result.

 FOR MORE VAGINITIS PANEL SAMPLE LAB REPORTS, [CLICK HERE](#)

